



Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170.
Controls Dandelion, Clover, Knotweed, Henbit, Chickweed, Plantain, Spurge and many other listed broadleaf weeds.

ACTIVE INGREDIENTS:

Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	30.56%
Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid	8.17%
Dimethylamine salt of dicamba: 3,6-dichloro-o-anisic acid	2.77%
INERT INGREDIENTS	58.50%

TOTAL	100.00%
--------------	----------------

THIS PRODUCT CONTAINS:

2.38 lb 2,4-dichlorophenoxyacetic acid equivalent per gallon or 25.38%
0.63 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per gallon or 6.75%
0.21 lb 3,6-dichloro-o-anisic acid equivalent per gallon or 2.30%
Isomer Specific by AOAC Methods.

KEEP OUT OF REACH OF CHILDREN DANGER-PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

—KEEP FROM FREEZING—

STOP! READ THE ENTIRE LABEL FIRST.

OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR ON CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment advice.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

SEE INSIDE LEAFLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND COMPLETE DIRECTIONS FOR USE.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Nonrefillable containers with capacities equal to or less than 5 gallons:

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Nonrefillable containers with capacities greater than 5 gallons: CONTAINER

HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

For Refillable Containers: CONTAINER HANDLING: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Container cleaning: Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

EPA Reg. No. 2217-656-5905

AD 090811

EPA Est. No. 2217-KS-1

NET CONTENTS: 2.5 Gallons (9.46 L)

Manufactured For

HELENA CHEMICAL COMPANY

225 SCHILLING BOULEVARD, SUITE 300
COLLIERVILLE, TENNESSEE 38017

PEEL BACK BOOK HERE AND RESEAL AFTER OPENING



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: Corrosive. Causes irreversible eye damage. Harmful if swallowed. Harmful if absorbed through skin. Do not get in eyes, on skin, or on clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are natural rubber. If you want more options, follow the instructions for Category A on an EPA chemical-resistance category selection chart.

All mixers, loaders, applicators and other handlers must wear:

- protective eyewear,
- long-sleeved shirt and long pants,
- shoes and socks,
- chemical-resistant gloves and
- chemical-resistant apron when mixing or loading, cleaning up spills or equipment, or otherwise exposed to the concentrate

See "Engineering Controls" for additional requirements.

ENGINEERING CONTROLS STATEMENTS

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

USER SAFETY RECOMMENDATIONS

- Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwaters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USES

Use this product only in accordance with its labeling.

This standard contains requirements for the protective handlers of agricultural pesticides. It contains requirements for the use of this product in accordance with the requirements of the Protection Standard.

Do not enter or allow worker entry into treated areas.

PPE required for early entry to treated areas that is with anything that has been treated, such as plants,

- coveralls worn over short-sleeved shirt and short pants,
- chemical-resistant footwear plus socks,
- chemical-resistant gloves made of any water-proof material,
- chemical-resistant headgear for overhead exposure,
- protective eyewear

NON-AGRICULTURAL USES

The requirements in this box apply to uses of this product that are not listed on the label. The WPS applies to agricultural pesticides (40 CFR Part 170). The WPS applies to forests, nurseries, or greenhouses.

Reentry Statement: Do not enter or allow people to enter treated areas until the drift has settled.

STORAGE

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a cool, dry place.

PESTICIDE DISPOSAL: Pesticide wastes are acutely toxic. Do not contaminate water, food, or feed by storage or disposal. A spill or release is a violation of Federal law. If these wastes cannot be disposed of by the manufacturer, they must be disposed of by a State Pesticide or Environmental Control Agency, or by other means approved by the local government.

Nonrefillable containers with capacities equal to or less than 5 gallons: Nonrefillable container. Do not reuse or refill. Dispose of in a sanitary landfill, or by incineration, or by other means approved by the local government.

Triple rinse as follows: Empty the remaining concentrate into the rinsate container. Fill the container 1/4 full with water or a mix tank or store rinsate for later use or disposal. Insert pressure rinsing nozzle and rinse for 10 seconds.

Triple rinse or pressure rinse container (or equivalent) and any remaining concentrate into the rinsate container.

Pressure rinse as follows: Empty the remaining concentrate into the rinsate container. Fill the container 1/4 full with water or a mix tank or store rinsate for later use or disposal. Insert pressure rinsing nozzle and rinse for 10 seconds.

Pressure rinse as follows: Empty the remaining concentrate into the rinsate container. Fill the container 1/4 full with water or a mix tank or store rinsate for later use or disposal. Insert pressure rinsing nozzle and rinse for 10 seconds.

ACTIONS CAREFULLY.

S
in. Do not get in eyes, on
ng gum, using tobacco or

ollow the instructions for

o the concentrate

orker Protection Standard
r modified as specified in

st, use detergent and hot
at have been drenched or

n clean clothing. If pes-
oving. As soon as pos-

apply directly to water, to
off may be hazardous to
nt washwaters or rinsate.
e of this chemical in areas
ation. Application around

. Only protected handlers

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170.

This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- coveralls worn over short-sleeved shirt and short pants,
- chemical-resistant footwear plus socks,
- chemical-resistant gloves made of any water-proof material,
- chemical-resistant headgear for overhead exposure
- protective eyewear

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Reentry Statement: Do not enter or allow people (or pets) to enter the treated area until sprays have dried.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

Nonrefillable containers with capacities equal to or less than 5 gallons:

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

(continued)

STORAGE

Nonrefillable containers with capacities greater than 5 gallons: **CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

For Refillable Containers:

CONTAINER HANDLING: Refillable container. Do not reuse or refill this container. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Container cleaning: Cleaning the container before reuse or before refilling is the responsibility of the user. To clean the container before final disposal, empty tank. Fill the container about 10 percent full with water or pump rinsate into application equipment or rinsate tank.

To clean the container before final disposal, empty tank. Fill the container about 10 percent full with water or pump rinsate into application equipment or rinsate tank.

SPRAYING

A variety of factors including weather conditions (temperature, wind speed, humidity, etc.) and ground application can influence pesticide drift. To minimize drift, apply this product.

Droplet Size

When applying sprays that contain 2,4-D mixed with a Medium or coarser spray (ASAE standard 572) or

Wind Speed

Do not apply at wind speeds greater than 10 mph. Do not apply in sensitive areas (including, but not limited to, residential areas, schools, hospitals, etc.) within 250 feet downwind. If applying a Medium spray, less than 10 mph.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator should be aware of atmospheric conditions exist at or below nozzle height. Do not apply in atmospheric conditions.

Susceptible Plants

Do not apply under circumstances where spray drift from the application may be expected to damage susceptible plants (including, but not limited to, growing stage), fruit trees (foliage), soybeans (vegetable), tobacco. Small amounts of spray drift that might occur are not considered a problem.

Other State and Local Requirements

Applicators must follow all State and local pesticide regulations, they must be observed.

Equipment

All ground application equipment must be properly calibrated and maintained. Nozzle height greater than 4 feet above the target area.



Part 170.
ries, and greenhouses, and
and emergency assistance.
personal protective equip-
t are covered by the Worker

urs.
d and that involves contact

ker Protection Standard for
gricultural plants on farms,
ve dried.

or pets. Keep from freezing.
icide, spray mixture, or rin-
l instructions, contact your
est EPA Regional Office for

ycling, if available, or punc-
by burning. If burned, stay

d drain for 10 seconds after
rsate into application equip-
ns to drip. Repeat this pro-

nk and continue to drain for
r mix tank or collect rinsate
t 40 PSI for at least 30 sec-

(continued)

STORAGE AND DISPOSAL (cont.)

Nonrefillable containers with capacities greater than 5 gallons:

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Triple rinse or pressure rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

For Refillable Containers:

CONTAINER HANDLING: Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose.

Container cleaning: Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of ground application can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

Droplet Size

When applying sprays that contain 2,4-D mixed with other active ingredients that require a Medium or more fine spray, apply only as a Medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Wind Speed

Do not apply at wind speeds greater than 10 mph. Only apply this product if the wind direction favors on-target deposition and there are not sensitive areas (including, but not limited to, bodies of water, known habitat for non-target species, nontarget crops) within 250 feet downwind. If applying a Medium spray, leave one swath unsprayed at the downwind edge of the treated field.

Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: a) conditions of temperature inversion exist, or b) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

Other State and Local Requirements

Applicators must follow all State and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

Equipment

All ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates. Do not apply with a nozzle height greater than 4 feet above the target site.

Product Description

Designed for turfgrass applications, non-crop, active ingredients.

- 2,4-D is an auxin-type herbicide which is a to the growing points of the plant, causing v
 - MCP-P (mecoprop-p) is an auxin-type her and is translocated to the growing points of tual plant death.
 - Dicamba is absorbed through the leaves an
- Combining these herbicides provides a very w affecting multiple sites within the broadleaf wee ing, and weed yellowing.

These combined herbicides provide limited res

EndRun® Offers These Advantages:

- Excellent postemergent activity with proven
- Extra 2,4-D for fast knockdown of dandelion
- The combinations of these active ingredients grass, including: dandelion, spurge and whit
- Rainfast in as little as 8 hours
- Controls major turf weeds
- Often, the weed injury symptoms can be not

Spray Preparation and Tank Mixes

In certain applications, liquid fertilizer may repli

Mixing with water:

Add one-half the required amount of water to tl with water. Mix thoroughly and continue agitati agitate to assure uniformity of the spray mixtur

Mixing with Liquid Fertilizers:

Use suitable sources and rates of fertilizer bas Verify physical compatibility with a jar test: Alw conducted by mixing all components in a small be mixed readily by shaking, then the mixture c flakes, sludge, gels or other precipitates form, i tilizer must not be prepared as a tank mixture.

Liquid fertilizers are either solutions (true fluids be satisfactory (may be marginal) without pre- ensure that the dispersants enable the herbicid

Adjuvants and Spray Additives:

Adjuvants (such as surfactants, spreaders, sp reducing agents) combined with this product (objectionable or would be unacceptable, then vants and spray additive tank-mix combination When an adjuvant is used with this product, Association (CPDA) certified adjuvant.

Ground Equipment

Spray distribution: The accuracy and unifor sprayers fitted with a boom or spray wand/g equipped with appropriate nozzles, tips, and s age, select a spray volume and delivery system Spray volumes of 5 to 220 gallons per acre wit greens) spray volumes, use 145 to 220 gallons per acre or 5 gallons per 1,000 square feet).

- Calibration and proper application are essen
- Overapplication or rates above those specifi
- Hand-held technique: Wands fitted with flat not be waved in a back-and-forth motion, or held stationary at the proper height. Side-to-



t.)

recycling, if available, or puncturing, by burning. If burned, stay

k. Fill the container 1/4 full with at least one complete revolution, then pour the contents over onto its other end and store for later use or

Use the tank and continue to drain for 1 minute or mix tank or collect rinsate about 40 PSI for at least 30 seconds.

Use this container for any other

When disposing of the container.

Use application equipment or a mix tank with the pump for 2 minutes. Pour the mixture and edure two more times.

relative humidity) and method of application. Make appropriate adjustments when

For a more fine spray, apply only as directed after for spinning atomizer nozzles.

For on-target deposition and there is no drift to nontarget crops (in the case of beans, and other vegetables, and other crops) within 100 feet of the treated field.

Temperature inversion exist, or b) stable temperature inversions or stable

that might be damaged or crops such as cotton, okra, flowers, grapes (in the case of beans, and other vegetables, and other crops) within 100 feet of the treated field.

-D herbicides. Where states have

substitutes or surrogates. Do not apply

Product Description

Designed for turfgrass applications, non-crop, and IVM (Industrial Vegetation Management) applications, this product contains three active ingredients.

1. 2,4-D is an auxin-type herbicide which is a class of plant growth regulators. It is absorbed through the leaves and is translocated to the growing points of the plant, causing weed stems curl and twist, leaf cupping and withering, and eventual plant death.
2. MCPP-p (mecoprop-p) is an auxin-type herbicide, which is a class of plant growth regulators. It is absorbed through the leaves and is translocated to the growing points of the plant, causing weed stems curl and twist, leaf cupping and withering, and eventual plant death.
3. Dicamba is absorbed through the leaves and roots and has multiple modes of action for hard-to-kill broadleaf weeds.

Combining these herbicides provides a very wide spectrum of weed control for susceptible weeds. **EndRun®** controls weeds by affecting multiple sites within the broadleaf weeds. The symptoms of susceptible broadleaf weeds include leaf and stem curl or twisting, and weed yellowing.

These combined herbicides provide limited residual activity.

EndRun® Offers These Advantages:

- Excellent postemergent activity with proven performance.
- Extra 2,4-D for fast knockdown of dandelions
- The combinations of these active ingredients provide effective weed control for common and troublesome weed species in turfgrass, including: dandelion, spurge and white clover.
- Rainfast in as little as 8 hours
- Controls major turf weeds
- Often, the weed injury symptoms can be noticed within hours of the application and plant death can occur within 14 to 21 days.

Spray Preparation and Tank Mixes

In certain applications, liquid fertilizer may replace part of the water.

Mixing with water:

Add one-half the required amount of water to the spray tank, then add **EndRun®** slowly with agitation, and complete filling the tank with water. Mix thoroughly and continue agitation while spraying. When this product is left standing for extended periods of time, re-agitate to assure uniformity of the spray mixture.

Mixing with Liquid Fertilizers:

Use suitable sources and rates of fertilizer based upon advice of your fertilizer supplier or State Extension Service Specialist.

Verify physical compatibility with a jar test: Always perform a jar test for compatibility before large scale mixing. The jar test can be conducted by mixing all components in a small container in proportionate quantities. If the mixture separates after standing and can be mixed readily by shaking, then the mixture can be used and applied with spray equipment providing continuous agitation. If large flakes, sludge, gels or other precipitates form, or if a separate oily layer or oil globules appear, then the herbicide and the liquid fertilizer must not be prepared as a tank mixture.

Liquid fertilizers are either solutions (true fluids) or suspensions. Mixing this product with suspensions or N-P-K solutions may not be satisfactory (may be marginal) without pre-mixing this product with water. Premixing this product with 2 to 4 parts water will ensure that the dispersants enable the herbicide to be suspended in the fertilizer.

Adjuvants and Spray Additives:

Adjuvants (such as surfactants, spreaders, spreader-stickers, spray thickeners, foaming agents, activators, detergents, and drift reducing agents) combined with this product can damage the leaf tissue of turfgrass. If any discoloration or cosmetic effects are objectionable or would be unacceptable, then adjuvant(s) combined with **EndRun®** would not be recommended. Do not use adjuvants and spray additive tank-mix combinations, unless your experience indicates that the tank mixture will not result in turf injury. When an adjuvant is used with this product, the manufacturer recommends the use of Chemical Producers and Distributors Association (CPDA) certified adjuvant.

Ground Equipment

Spray distribution: The accuracy and uniformity of the herbicide distribution is the sole responsibility of the applicator. Power sprayers fitted with a boom or spray wand/gun may be used for broadcast applications and spot treatments. Boom sprayers equipped with appropriate nozzles, tips, and screens are suitable for broadcast applications. For best spray distribution and coverage, select a spray volume and delivery system that will ensure accurate and uniform coverage.

Spray volumes of 5 to 220 gallons per acre with spray pressures adjusted to between 20 to 40 psi. **Note:** For bentgrass (except golf greens) spray volumes, use 145 to 220 gallons per acre. Use higher spray volumes for dense weed populations (up to 220 gallons per acre or 5 gallons per 1,000 square feet).

- Calibration and proper application are essential when using this product.
- Overapplication or rates above those specified on this label can cause turf injury.
- Hand-held technique: Wands fitted with flat fan nozzle tips may be used with the appropriate technique. Flat fan nozzles should not be waved in a back-and-forth motion, or in a side-to-side motion, or in a swinging arm motion. Instead, the nozzle should be held stationary at the proper height. Side-to-side motion results in uneven coverage.

Hand-operated sprayers including backpack sprayers. After using this product, clean sprayer with water before applying other pesticides.

Where to Use

This product provides broadleaf control in turfgrass.

Ornamental Turfgrass Sites:

- **Residential/domestic sites** are defined as lawns, apartment complexes, condominiums, and patios.
- **Ornamental turf sites** include turfgrass in institutional buildings, recreation areas, fairways, and golf courses.
- **Institutional sites** are defined as turf areas including, but not limited to hospitals, nursing homes, and office buildings.
- **Non-cropland sites:** including farmyard and unpaved roads; Roadsides, roadside ditches, Lumberyards, tank farms, fuel or equipment storage areas, railroad rights-of-ways, railroad yards, rail yards, and communication lines.
- **Agricultural sites:** Commercial sod production.

Prohibitions of Sites:

- Do not apply to any body of water such as any shorelines (non-cropland sites adjacent to saltwater bays).
- Do not apply to wetlands (swamps, bogs, and marshes).
- Do not apply to agricultural irrigation water.
- Do not apply to agricultural drainage water.
- Applications to non-cropland areas are not recommended for commercial use, or for commercial use.
- Do not apply or allow this product to come in contact with ornamental trees, or other desirable broadleaf trees or shrubs.

Turfgrass Tolerance:

- The turfgrass tolerance to this product may vary due to the selectivity on the turfgrass.
- Certain spray tank additives (adjuvants, surfactants, and concentrates) may reduce the selectivity of this product. Your experience indicates that the tank mixture manufacturer recommends the use of Chemical Producers and Distributors Association (CPDA) certified adjuvant.

Prohibitions:

- Do not apply this product to St. Augustine grass where St. Augustine grass is present.
- Do not broadcast apply this product where air temperatures exceed 90°F. Where State regulations must be observed.
- To avoid turf injury, use only on turfgrass where excess rainfall/irrigation, shaded areas, or other conditions may occur if this product is applied under conditions that may cause damage caused by the use of this product.
- For ground application only; aerial application is not recommended.
- Do not use this product on or near desirable trees or shrubs within the dripline of desirable trees and shrubs.

Application Schedules

Apply this product to broadleaf weeds that are present in turfgrass areas and provide control for emerged winter annuals and perennials.

ications, this product contains three
rough the leaves and is translocated
ering, and eventual plant death.
rs. It is absorbed through the leaves
af cupping and withering, and even-
d-to-kill broadleaf weeds.
weeds. **EndRun®** controls weeds by
Js include leaf and stem curl or twist-

d troublesome weed species in turf-

path can occur within 14 to 21 days.

gitation, and complete filling the tank
ing for extended periods of time, re-

Extension Service Specialist.
rge scale mixing. The jar test can be
ure separates after standing and can
roviding continuous agitation. If large
then the herbicide and the liquid fer-

ensions or N-P-K solutions may not
product with 2 to 4 parts water will

nts, activators, detergents, and drift
discoloration or cosmetic effects are
be recommended. Do not use adju-
k mixture will not result in turf injury.
hemical Producers and Distributors

sponsibility of the applicator. Power
nd spot treatments. Boom sprayers
or best spray distribution and cover-
.

osi. **Note:** For bentgrass (except golf
weed populations (up to 220 gallons

e technique. Flat fan nozzles should
otion. Instead, the nozzle should be

Hand-operated sprayers including backpack sprayers, compression sprayers are appropriate for small turfgrass areas.

After using this product, clean sprayer with soap or detergent and water, or an approved spray tank cleaner and rinse thoroughly before applying other pesticides.

Where to Use

This product provides broadleaf control in the following sites:

Ornamental Turfgrass Sites:

- **Residential/domestic sites** are defined as areas associated with the household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes.
- **Ornamental turf sites** include turfgrass established around residences, parks, streets, retail outlets, cemeteries, industrial and institutional buildings, recreation areas, fairgrounds, areas adjacent to athletic fields and paved areas.
- **Institutional sites** are defined as turf areas around properties or facilities providing a service to public or private organizations including, but not limited to hospitals, nursing homes, schools, museums, libraries, sport facilities, golf courses (fairways and roughs), and office buildings.
- **Non-cropland sites:** including farmyards, fencerows or fence lines, highway rights-of-way (principal, interstate, county, private, and unpaved roads): Roadsides, roadside ditches, road shoulders, road embankments, dividers, and medians; Industrial sites: Lumberyards, tank farms, fuel or equipment storage areas; Municipal, State, and Federal lands: Airports and military installations; railroad rights-of-ways, railroad yards, railroad crossings and railroad bridge abutments; Utility rights-of-way: telephone, pipeline, electrical powerlines, and communication transmission lines.
- **Agricultural sites:** Commercial sod production

Prohibitions of Sites:

- Do not apply to any body of water such as lakes, streams, rivers, ponds, reservoirs, or estuaries (saltwater bays). Do not apply to any shorelines (non-cropland sites adjacent to the edges of a body of water) for lakes, streams, rivers, ponds, reservoirs, or estuaries (saltwater bays).
- Do not apply to wetlands (swamps, bogs, potholes, or marshes).
- Do not apply to agricultural irrigation water or on agricultural irrigation ditchbanks and canals.
- Do not apply to agricultural drainage water or on agricultural ditchbanks.
- Applications to non-cropland areas are not applicable to treatment of commercial timber or other plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.
- Do not apply or allow this product to come into direct contact with cotton, grapes, tobacco, vegetable crops, flowers, fruit or ornamental trees, or other desirable broadleaf plants; small amounts of spray drift may injure susceptible plants, including ornamental trees or shrubs.

Turfgrass Tolerance:

- The turfgrass tolerance to this product may vary and temporary turfgrass yellowing may occur. Adverse environmental conditions may reduce the selectivity on the turfgrass. Do not apply this product to stressed turf.
- Certain spray tank additives (adjuvants, wetting agents, surfactants), liquid fertilizers, and tank mixtures containing emulsifiable concentrates may reduce the selectivity on the turfgrass. Do not use adjuvants and spray additive tank-mix combinations, unless your experience indicates that the tank mixture will not result in turf injury. When an adjuvant is used with this product, the manufacturer recommends the use of Chemical Producers and Distributors Association (CPDA) certified adjuvant.

Prohibitions:

- Do not apply this product to St. Augustinegrass, bentgrass greens, carpetgrass, dichondra, legumes, and lawns where desirable clovers are present.
- Do not broadcast apply this product when temperatures are above 90°F, some injury may be expected with spot treatments when air temperatures exceed 90°F. Where State, county or local governments have more stringent temperature regulations, these regulations must be observed.
- To avoid turf injury, use only on turfgrass that is reasonably free of stress from diseases, insects, excess heat or cold, drought or excess rainfall/irrigation, shaded areas, nematodes, improper mowing or improper applications of fertilizer and pesticides. Injury can occur if this product is applied under any of these or other stress conditions. Under any of these stress conditions, any turf damage caused by the use of this product is beyond the control of the registrant and all risk is assumed by the buyer and/or user.
- For ground application only; aerial applications are not permitted.
- Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the dripline of desirable trees and shrubs, since injury may result.

Application Schedules

Apply this product to broadleaf weeds that are young and actively growing for the best results. Fall applications can provide improved control for emerged winter annuals and perennials such as henbit, chickweed, clover and ground ivy.



For the Listed Residential/domestic sites, Ornamental Turf sites, Institutional sites and Agricultural sites:

Do not apply more than two (2) broadcast treatments of this product per site per year. A second broadcast application or a follow-up application as a spot treatment is advised for more mature weeds, for dense infestations, and for adverse environmental conditions.

Spot treatments during the summer may be appropriate for sparse infestations, or as a follow-up treatment, or any time broadleaf weeds are actively growing.

For the Listed Non-cropland sites:

Use only one (a single) broadcast treatment of this product per site per year for woody plant infestations, or two broadcast treatments for annual and perennial weeds. It is required to wait 30 days between treatments and a spot treatment may be substituted for the second broadcast treatment if necessary.

Extremes in environmental conditions (e.g., temperature and moisture, soil conditions, and cultural practices) may affect the activity of this product. Under warm moist conditions, herbicide symptoms may be accelerated. While under very dry conditions, the expression of herbicide symptoms is delayed, and weeds hardened off by drought are less susceptible to this product.

For newly seeded areas:

Delay application of this product to grass seedlings until after the second mowing.

For newly sodded, sprigged, or plugged areas:

The application of this product to newly sodded, sprigged, or plugged grasses should be delayed until 3 to 4 weeks after the sodding, sprigging, or plugging operations.

Reseeding interval:

Treated areas may be reseeded 3 weeks after application.

How Much to Use**USE RATES AND SPRAY VOLUMES:**

Generally, the lower application rates within the specified range will provide satisfactory control of sensitive weed species. The higher application rates within the specified range will be required for dense infestations of perennial weeds, for adverse/extreme environmental conditions, or for weeds hardened off or more mature.

TABLE 1. USE RATES FOR ORNAMENTAL TURF SITES, NON-CROPLAND AND SOD FARMS		
Species	Rate	Spray Volume
Cool-Season Turf		
Kentucky bluegrass, annual bluegrass, annual ryegrass, perennial ryegrass, tall fescue, red or fine leaf fescues	3.0 to 4 Pints/Acre (1.1 to 1.5 fl oz/1,000 sq ft)	5 to 220 Gallons/Acre (0.125 to 5.0 Gallons/1,000 sq ft)
Creeping Bentgrass (excluding golf greens) Apply preferably in May or mid-August through September. Slight turf yellowing will disappear after about one (1) week. Note: Care should be taken to avoid overdosing bentgrass or injury may result.	1.8 Pints/Acre (0.66 fl oz/1,000 sq ft)	145 to 220 Gallons/Acre (3.33 to 5.0 Gallons/1,000 sq ft) High spray volumes will provide uniform coverage.
Warm-Season Turf		
Hybrid bermudagrass, common bermudagrass, zoysiagrass, buffalograss and bahiagrass Slight turf yellowing will disappear after about one (1) week.	3.0 to 4 Pints/Acre (1.1 to 1.5 fl oz/1,000 sq ft) Sod Farms: 2 to 2.25 Pints/Acre (0.75 to 0.83 fl oz/1,000 sq ft)	5 to 220 Gallons/Acre (0.125 to 5.0 Gallons/1,000 sq ft)
Note: Do not apply to above listed warm-season turfgrass unless some turf injury can be tolerated. It is impossible to test all environmental conditions for the listed warm-season turfgrass. Do not apply this product to warm-season turfgrass during spring green-up or in the fall during the transition period between active growth and dormancy. Dormant turf: This product may be applied to fully dormant bermudagrass, fully dormant Zoysiagrass, fully dormant buffalograss and fully dormant bahiagrass.		

SPOT TREATMENT:**WITH HAND OPERATED SPRAYERS (INCLUDING BROADCASTERS):**

- Apply any time the emerged broadleaf weeds are active.
- Calibration and proper application are essential when using this product.
- Uniform applications are essential when using this product. Excessive overlaps of this product can cause turf injury.
- Hand-held techniques: Wands fitted with flat fan nozzle not be waved in a back-and-forth motion, or in a side-to-side motion, or in a side-to-side motion. Side-to-side motion held stationary at the proper height. Side-to-side motion.
- Follow-up applications as spot treatments at a 30-day adverse environmental conditions.
- **For cool-season turfgrass (except Bentgrass) list water for treatment of approximately 1,000 sq ft of turf.**
- **For Bentgrass (excluding golf greens):** Mix 0.66 fl oz/1,000 sq ft of turfgrass. This high spray volume is susceptible.
- **For warm-season turfgrass listed in Table 1:** Mix of approximately 1,000 sq ft of turfgrass. Apply any time.
- **Limitations on spot treatments to residential turf:** The maximum application rate is 1.5 fl oz/1,000 sq ft per acre. The maximum number of spot treatments is limited to 10 per acre.

CULTURAL TIPS:**Irrigation:**

- Do not apply this product through any type of irrigation.
- Rainfall in as little as 8 hours. Do not apply this product if rainfall is less than 8 hours.
- If possible, do not irrigate or water the turfgrass within 48 hours of application.
- If dry conditions exist, a scheduled irrigation or watering is recommended.

Mowing:

For optimum results, delay mowing 2 days before and after application.

Broadleaf Weeds Controlled

EndRun® will control the following broadleaf weeds. Apply as directed.

BROADLEAF WEEDS	
annual fleabane aster, white heath & white prairie bedstraw beggarticks beggarweed, creeping bindweed birdsfoot trefoil black medic broadleaf plantain buckhorn plantain bull thistle burdock burdock, common Carolina geranium carpetweed chickweed, common chicory cinquefoil clover cocklebur compassplant curly dock dandelion dayflower deadnettle	dock dogfennel dovefoot geranium false dandelion (*spotted catsear & common) field bindweed (*morningglory & creeping) field madder field oxeye-daisy (*creeping) field pennycress filaree, whitestem & redstem Florida pusley ground ivy groundsel hairy bittercress hawkweed healall henbit horsenettle horsetweed innocence (Blue-eyed Mary) jimsonweed kochia knotweed lamb'squarters
*Synonyms **A repeat application may be required in 30 days.	

tural sites:

application or a follow-up environmental conditions.

or, any time broadleaf

or two broadcast treatment may be substituted

) may affect the activity conditions, the expression.

4 weeks after the sod-

eed species. The higher volume/extreme environ-

ly Volume
Gallons/Acre
Gallons/1,000 sq ft)
0 Gallons/Acre
Gallons/1,000 sq ft)
olumes will provide m coverage.
Gallons/Acre
Gallons/1,000 sq ft)
ossible to test all envi-
sition period between
fully dormant buffalo-

SPOT TREATMENT:**WITH HAND OPERATED SPRAYERS (INCLUDING BACKPACK SPRAYERS AND PUMP-UP TYPE SPRAYERS):**

- Apply any time the emerged broadleaf weeds are actively growing.
- Calibration and proper application are essential when using this product.
- Uniform applications are essential when using this product. Overapplication or rates above those specified on this label including excessive overlaps of this product can cause turf injury.
- Hand-held techniques: Wands fitted with flat fan nozzle tips may be used with the appropriate technique. Flat fan nozzles should not be waved in a back-and-forth motion, or in a side-to-side motion, or in a swinging arm motion. Instead, the nozzle should be held stationary at the proper height. Side-to-side motion results in uneven coverage.
- Follow-up applications as spot treatments at a 30-day interval are suitable for more mature weeds, for dense infestations, and for adverse environmental conditions.
- **For cool-season turfgrass (except Bentgrass) listed in Table 1:** Mix 1.1 to 1.5 fl oz of this product per one (1.0) gallon of water for treatment of approximately 1,000 sq ft of turfgrass. Apply any time the emerged broadleaf weeds are susceptible.
- **For Bentgrass (excluding golf greens):** Mix 0.66 fl oz of this product per four (4.0) gallons of water for treatment of approximately 1,000 sq ft of turfgrass. This high spray volume will provide uniform coverage. Apply any time the emerged broadleaf weeds are susceptible.
- **For warm-season turfgrass listed in Table 1:** Mix 0.75 to 0.83 fl oz of this product per one (1.0) gallon of water for treatment of approximately 1,000 sq ft of turfgrass. Apply any time the emerged broadleaf weeds are susceptible.
- **Limitations on spot treatments to residential turfgrass:** Spot treatment is defined as a treatment area no greater than 1,000 sq ft per acre. The maximum application rate is 1.5 fl oz per 1,000 sq ft per application (0.33 lb MCP-P acid equivalent per acre). The maximum number of spot treatments is limited to 2 per year with a minimum of 30 days between applications.

CULTURAL TIPS:**Irrigation:**

- Do not apply this product through any type of irrigation system.
- Rainfall in as little as 8 hours. Do not apply this product immediately before rainfall or irrigation.
- If possible, do not irrigate or water the turfgrass within 8 to 24 hours after application.
- If dry conditions exist, a scheduled irrigation or watering 24 hours before and 24 hours after application is recommended.

Mowing:

For optimum results, delay mowing 2 days before and until 2 days after the application of this product.

Broadleaf Weeds Controlled

EndRun® will control the following broadleaf weeds. Apply any time the emerged broadleaf weeds are susceptible.

BROADLEAF WEEDS			
annual fleabane	dock	lawn burweed	roundleaf greenbrier
aster, white heath & white prairie	dogfennel	lespedeza, common	shepherdspurge
bedstraw	dovefoot geranium	mallow, common	spotted spurge
beggarticks	false dandelion	matchweed	spurge
beggarweed, creeping	(*spotted catsear & common catsear)	mouseear chickweed	sunflower
bindweed	field bindweed	mustard	thistle
birdsfoot trefoil	(*morningglory & creeping jenny)	nettle	velvetleaf (*buttonweed)
black medic	field madder	old world diamond flower	Venice mallow
broadleaf plantain	field oxeye-daisy (*creeping oxeye)	<i>Oxalis</i> (*yellow woodsorrel)	<i>Veronica</i> (*corn speedwell)
buckhorn plantain	field pennycress	parsley-piert	Virginia buttonweed**
bull thistle	filaree, whitestem & redstem	Pennsylvania smartweed	Virginia-creeper
burclover	Florida pusley	pennycwort (*dollarweed)	western salsify
burdock, common	ground ivy	pepperweed	white clover (*Dutch clover,
Carolina geranium	groundsel	pigweed	honeyuckle clover,
carpetweed	hairy bittercress	pineappleweed	white trefoil & purplewort)
chickweed, common	hawkweed	plantain	wild carrot
chicory	healall	poison ivy	wild garlic
cinqfoil	henbit	poison oak	wild geranium
clover	horsenettle	prostrate knotweed (*knotweed)	wild lettuce
cocklebur	horsetweed	puncturevine	wild mustard
compassplant	innocence (Blue-eyed Mary)	purple cudweed	wild onion
curly dock	jimsonweed	purslane	wild strawberry
dandelion	kochia	ragweed	yarrow
dayflower	knotweed	redweed	yellow rocket
deadnettle	lambsquartars	red sorrel (*sheep sorrel)	and many more broadleaf weeds

*Synonyms **A repeat application may be required in 30 days.

For Use in Non-Cropland: Brush and Weed Control High volume foliar applications:

Apply up to 1.15 gallons of product per 100 gallons of high volume equipment. Use the lower spray concentrations within the range for hard-to-control species, conditions (e.g., drought).

Spray broadleaf weeds, woody plants or mixed brush. The minimum total volume of spray solution required for tion per treated acre. The spray preparation chart for

TABLE 3. Instructions for preparing 25 to 100 gallon volume foliar applications.

Spray solution per acre, Gallons	
25	
75	
100	

Equal measures: 1 gallon = 4 quarts = 8 pints = 128

For Backpack Sprayers and Hand Pump-up Spr

TABLE 4. Instructions for preparing 1 to 3 gallons c ume foliar applications.

Gallons of Water	
1	
2	
3	

Equal measures: 1 fl oz = 2 tablespoons (Tbs) = 6 te

BRUSH

Ash	Cedars
Aspen	Elms
Birch	Gooseberry
Black cherry	Honey locust
Brambles	Kudzu

TABLE 2. Restrictions on broadcast applications to

Use Site	Maximum Rate per Application
Ornamental turfgrass sites and sod farms	4.0 pints/A (1.19 lb 2,4-D ae/A) (0.33 lb MCP-P ae/A) (0.11 lb dicamba ae/A)

**SPRAYERS):**

fied on this label including

je. Flat fan nozzles should
stead, the nozzle should be
dense infestations, and for

luct per one (1.0) gallon of
eds are susceptible.
r for treatment of approxi-
emerged broadleaf weeds

illon of water for treatment
area no greater than 1,000
> acid equivalent per acre).
applications.

is recommended.

usceptible.

leaf greenbriar
herdspurse
ed spurge
je
ower
e
tleaf (*buttonweed)
e mallow
nica (*corn speedwell)
ia buttonweed**
ia-creeper
ern salsify
> clover (*Dutch clover,
neysuckle clover,
ite trefoil & purplewort)
carrot
garlic
geranium
lettuce
mustard
onion
strawberry
w
w rocket
many more broadleaf weeds

For Use in Non-Cropland: Brush and Weed Control**High volume foliar applications:**

Apply up to 1.15 gallons of product per 100 gallons of water or apply a 1.0 to 1.15% vol/vol spray solution as a full cover spray with high volume equipment. Use the lower spray concentrations in the range for susceptible species and use the higher spray concentrations within the range for hard-to-control species, for mature plants during the late summer or under adverse environmental conditions (e.g., drought).

Spray broadleaf weeds, woody plants or mixed brush uniformly and thoroughly by wetting all leaves, stems, bark and root collars. The minimum total volume of spray solution required for adequate coverage of solid stands of mixed brush is 25 gallons of spray solution per treated acre. The spray preparation chart for applications on a spray-to-wet basis is shown below in **Table 3**.

TABLE 3. Instructions for preparing 25 to 100 gallons of spray solution at 1.0 to 1.15% spray concentration with water for high volume foliar applications.

Spray solution per acre, Gallons	Amount of Product Needed for Spray Concentration of:	
	1.0%	1.15%
25	2.0 pt	2.3 pt
75	6.0 pt	6.9 pt
100	1.0 gal	1.15 gal

Equal measures: 1 gallon = 4 quarts = 8 pints = 128 fl oz

For Backpack Sprayers and Hand Pump-up Sprayers**TABLE 4.** Instructions for preparing 1 to 3 gallons of spray solution at 1.0 to 1.15% spray concentration with water for high volume foliar applications.

Gallons of Water	Amount of Product Needed for Spray Concentration of:	
	1.0%	1.15%
1	1.25 fl oz (2.5 Tablespoons)	1.5 fl oz (3 Tablespoons)
2	2.5 fl oz	3.0 fl oz
3	3.8 fl oz	4.5 fl oz

Equal measures: 1 fl oz = 2 tablespoons (Tbs) = 6 teaspoons (tsp)

BRUSH

Ash	Cedars	Multiflora rose	Sumac
Aspen	Elms	Oak	Willow
Birch	Gooseberry	Poison ivy	And many more brush species
Black cherry	Honey locust	Poison oak	
Brambles	Kudzu	Shortleaf pine	

TABLE 2. Restrictions on broadcast applications to ornamental turfgrass sites, sod farms, and non-cropland.

Use Site	Maximum Rate per Application	Maximum Number of Applications per Year	Minimum Interval Between Applications	Maximum Seasonal Rate
Ornamental turfgrass sites and sod farms	4.0 pints/A (1.19 lb 2,4-D ae/A) (0.33 lb MCPP-p ae/A) (0.11 lb dicamba ae/A)	2	30 days	8.0 pints/A (2.37 lb 2,4-D ae/A) (0.65 lb MCPP-p ae/A) (0.22 lb dicamba ae/A)

(continued)

TABLE 2. Restrictions on broadcast application:

Use Site	Maximum Rate per Application
Non-cropland	4.0 pints/A (1.19 lb 2,4-D ae/A) (0.33 lb MCPP-p ae/A) (0.11 lb dicamba ae/A)

Use of this product in certain portions of California
relief in *Washington Toxics Coalition*, et.al. v. EPA
<http://www.epa.gov/espp/litstatus/wtc/qs-as.htm>.

LIMITED W

IMPORTANT: Read this LIMITED WARRANTY. At product, buyer and all users agree to accept the t exception. If the terms are not acceptable, return t will be refunded in full.

It is impossible to eliminate all risks inherently ass other unintended consequences can result from u materials, or the manner of use or application, etc **PURCHASING AND USING THIS PRODUCT TI SUCH RISKS.** Buyer and all users further agree to is not explicitly set forth in or that is inconsistent v

The manufacturer warrants only that this product c sonably suited for the labeled use when applied a **TO THE EXTENT CONSISTENT WITH APPLIC OTHER EXPRESS OR IMPLIED WARRANTIES A PARTICULAR PURPOSE, WHICH ARE HERI**

THE EXCLUSIVE REMEDY OF BUYER AND A MANUFACTURER, FOR ANY AND ALL LOSSE OF THIS PRODUCT, WHETHER OR NOT BASI WISE, SHALL BE LIMITED, AT THE MANUFAI PURCHASE PRICE FOR THE QUANTITY OF F EXTENT CONSISTENT WITH APPLICABLE LA TAL, CONSEQUENTIAL, OR SPECIAL DAMA Manufacturer must be promptly notified in writing wise, to be eligible to receive either remedy stated

The terms of this **LIMITED WARRANTY AND DI** the point of sale or elsewhere. No employee or ag Limited Warranty and Disclaimer in any manner.

EndRun® is a registered trademark of Helena Hol



tion as a full cover spray with use the higher spray concentration to control adverse environmental conditions.

stems, bark and root collars. Use is 25 gallons of spray solution in **Table 3**.

concentration with water for high

concentration of:

1.15%

2.3 pt

6.9 pt

1.15 gal

ation with water for high vol-

concentration of:

1.15%

1.5 fl oz
(3 Tablespoons)

3.0 fl oz

4.5 fl oz

mac
illow
and many more brush species

cropland.

Maximum Seasonal Rate
8.0 pints/A
(2.37 lb 2,4-D ae/A) (0.65 lb MCP-p ae/A) (0.22 lb dicamba ae/A)

(continued)

TABLE 2. Restrictions on broadcast applications to ornamental turfgrass sites, sod farms, and non-cropland. (continued)

Use Site	Maximum Rate per Application	Maximum Number of Applications per Year	Minimum Interval Between Applications	Maximum Seasonal Rate
Non-cropland	4.0 pints/A (1.19 lb 2,4-D ae/A) (0.33 lb MCP-p ae/A) (0.11 lb dicamba ae/A)	2 (annual and perennial weeds)	30 days	8.0 pints/A (2.37 lb 2,4-D ae/A) (0.65 lb MCP-p ae/A) (0.22 lb dicamba ae/A)

Use of this product in certain portions of California, Oregon, and Washington is subject to the January 22, 2004 Order for injunctive relief in *Washington Toxics Coalition, et.al. v. EPA*, C01-0132C, (W.D. WA). For further information, please refer to EPA web site: <http://www.epa.gov/espp/litstatus/wtc/qs-as.htm>.

LIMITED WARRANTY AND DISCLAIMER

IMPORTANT: Read this LIMITED WARRANTY AND DISCLAIMER before buying or using this product. By opening and using this product, buyer and all users agree to accept the terms of this LIMITED WARRANTY AND DISCLAIMER in their entirety and without exception. If the terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full.

It is impossible to eliminate all risks inherently associated with use of this product. Damage to the treated article, ineffectiveness, or other unintended consequences can result from use of the product under abnormal conditions such as weather, presence of other materials, or the manner of use or application, etc. Such factors and conditions are beyond the control of the manufacturer, and **BY PURCHASING AND USING THIS PRODUCT THE BUYER AND ALL USERS OF THIS PRODUCT AGREE TO ACCEPT ALL SUCH RISKS.** Buyer and all users further agree to assume all risks of loss or damage from the use of the product in any manner that is not explicitly set forth in or that is inconsistent with label instructions, warnings and cautions.

The manufacturer warrants only that this product conforms to the chemical description given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use, subject to the inherent risks described below. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED.**

THE EXCLUSIVE REMEDY OF BUYER AND ALL USERS OF THIS PRODUCT, AND THE EXCLUSIVE LIABILITY OF THE MANUFACTURER, FOR ANY AND ALL LOSSES, DAMAGES, OR INJURIES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER OR NOT BASED IN CONTRACT, NEGLIGENCE, STRICT LIABILITY IN TORT OR OTHERWISE, SHALL BE LIMITED, AT THE MANUFACTURER'S OPTION, TO REPLACEMENT OF OR THE REPAYMENT OF THE PURCHASE PRICE FOR THE QUANTITY OF PRODUCT WITH RESPECT TO WHICH DAMAGES ARE CLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. The Manufacturer must be promptly notified in writing of any claims, whether based in contract, tort, negligence, strict liability, or otherwise, to be eligible to receive either remedy stated above.

The terms of this **LIMITED WARRANTY AND DISCLAIMER** cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere. No employee or agent of the manufacturer or seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

EndRun® is a registered trademark of Helena Holding Company.

Use this product only in accordance with the label. Do not use on Dandelion, Clover, Knotweed, or other broadleaf weeds.

ACTIVE INGREDIENTS:

Dimethylamine salt of 2,4-dichlorophenoxyacetic acid
Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy)acetic acid
Dimethylamine salt of dicamba: 3,6-dichloro-2-methylbenzoic acid

INERT INGREDIENTS

TOTAL

THIS PRODUCT CONTAINS:

2.38 lb 2,4-dichlorophenoxyacetic acid equivalent
0.63 lb (+)-(R)-2-(2-methyl-4-chlorophenoxy)acetic acid equivalent
0.21 lb 3,6-dichloro-o-anisic acid equivalent per isomer Specific by AOAC Methods.

KEEP OUT OF REACH OF CHILDREN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

STOP! READ THE ENTIRE LABEL FIRST

IF IN EYES:	- Hold eye open and rinse slowly and gently with water for 15-20 minutes. - Remove contact lenses, if present, after the first irrigation. - Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	- Call a poison control center or doctor for treatment advice. - Have person sip a glass of water if able to swallow. - Do not induce vomiting unless told to do so by a poison control center or doctor. - Do not give anything by mouth to an unconscious person.
IF ON SKIN OR ON CLOTHING:	- Take off contaminated clothing. - Rinse skin immediately with plenty of water. - Call a poison control center or doctor for treatment advice.
IF INHALED:	- Move person to fresh air. - If person is not breathing, call 911 or doctor. - Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a doctor or emergency medical treatment advice.

NOTE TO PHYSICIAN: Probable mucosal damage may occur. Use special precautions and treatment.

SEE INSIDE LEAFLET FOR ADDITIONAL INFORMATION

AD 090811

EPA Reg. No. 2217-656-5905

HELENA
225 SCHILLING BOULEVARD

Job #57588

non-cropland. (continued)	
val	Maximum Seasonal Rate
	8.0 pints/A (2.37 lb 2,4-D ae/A) (0.65 lb MCPP-p ae/A) (0.22 lb dicamba ae/A)

January 22, 2004 Order for injunctive
relief, please refer to EPA web site:

R

product. By opening and using this
product in their entirety and without
purchase, and the purchase price

treated article, ineffectiveness, or
such as weather, presence of other
control of the manufacturer, and **BY
YOUR PURCHASE YOU AGREE TO ACCEPT ALL
RISKS** of the product in any manner that

label, and that the product is re-
sponsible for inherent risks described below.
**HELENA MAKES NO REPRESENTATIONS AND
WARRANTS NO MERCHANTABILITY OR FITNESS FOR**

**EXCLUSIVE LIABILITY OF THE
MANUFACTURER FROM THE USE OR HANDLING
OF THIS PRODUCT. NO OTHER
LIABILITY IN TORT OR OTHERWISE
OR THE REPAYMENT OF THE
DAMAGES ARE CLAIMED. TO THE
EXTENT PERMITTED BY LAW, THE
MANUFACTURER SHALL NOT BE LIABLE FOR INCIDENTAL
DAMAGES OF THIS PRODUCT.** The
negligence, strict liability, or other-

verbal statements or agreements at
no time vary or exceed the terms of this



**Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170.
Controls Dandelion, Clover, Knotweed, Henbit, Chickweed, Plantain, Spurge and many other listed broadleaf weeds.**

ACTIVE INGREDIENTS:

Dimethylamine salt of 2,4-dichlorophenoxyacetic acid	30.56%
Dimethylamine salt of (+)-(R)-2-(2-methyl-4-chlorophenoxy) propionic acid	8.17%
Dimethylamine salt of dicamba: 3,6-dichloro-o-anisic acid	2.77%

INERT INGREDIENTS

TOTAL	58.50%
	100.00%

THIS PRODUCT CONTAINS:

2.38 lb 2,4-dichlorophenoxyacetic acid equivalent per gallon or 25.38%
0.63 lb (+)-(R)- 2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per gallon or 6.75%
0.21 lb 3,6-dichloro-o-anisic acid equivalent per gallon or 2.30%
Isomer Specific by AOAC Methods.

KEEP OUT OF REACH OF CHILDREN DANGER – PELIGRO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

— KEEP FROM FREEZING —

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

FIRST AID

IF IN EYES:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF ON SKIN OR ON CLOTHING:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
IF INHALED:	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment advice.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

SEE INSIDE LEAFLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND COMPLETE DIRECTIONS FOR USE.

AD 090811

EPA Reg. No. 2217-656-5905

EPA Est. No. 2217-KS-1

NET CONTENTS: 2.5 Gallons (9.46 L)

Manufactured For
HELENA CHEMICAL COMPANY
225 SCHILLING BOULEVARD, SUITE 300 • COLLIERVILLE, TN 38017

Job #57588