

# FYFANON<sup>®</sup>

## Malathion Insecticide

**ACTIVE INGREDIENT:**

Malathion: 0,0-dimethyl phosphorodithioate of diethyl mercaptosuccinate..... 56.44%

**INERT INGREDIENTS** ..... 43.56%

**TOTAL** ..... 100.00%

(ONE GALLON OF FYFANON CONTAINS 5 POUNDS OF MALATHION.)

### KEEP OUT OF REACH OF CHILDREN WARNING – AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you.)

#### PRECAUTIONARY STATEMENTS Hazards to Humans and Domestic Animals WARNING – AVISO

Causes substantial but temporary eye injury. Harmful if swallowed. Harmful if absorbed through skin. Harmful if inhaled. Do not get in eyes or on clothing. Avoid breathing vapor or spray mist.

#### FIRST AID

<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"><li>• Immediately call a poison control center or doctor.</li><li>• Do not induce vomiting unless told to by a poison control center or doctor.</li><li>• Do not give <b>any</b> liquid to the person.</li><li>• Do not give anything by mouth to an unconscious person.</li></ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"><li>• Move person to fresh air.</li><li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li><li>• Call a poison control center or doctor for further treatment advice.</li></ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"><li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"><li>• Take off contaminated clothing.</li><li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>• Call a poison control center or doctor for treatment advice.</li></ul>

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency, call 1-800-424-9300.

**NOTE TO PHYSICIAN:** Contains an organophosphate that inhibits cholinesterase. Atropine is antidotal. 2PAM may be effective as an adjunct to atropine.

#### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

**PESTICIDE STORAGE:** FYFANON<sup>®</sup> should be stored in the original unopened container in a secure, dry place. Do not contaminate with other pesticides or fertilizers. The product should never be heated above 55°C (131°F), and should not be stored for long periods of time at a temperature in excess of 25°C (77°F).

**PESTICIDE DISPOSAL:** To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by State or local governments or by industry).

**CONTAINER DISPOSAL: Nonrefillable container (equal to or less than 5 gallons):** Do not reuse or refill this container. Triple rinse (or equivalent). Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

**Nonrefillable container (greater than 5 gallons):** Do not reuse or refill this container. Triple rinse (or equivalent). Triple rinse as follows: Empty the remaining contents into application equipment or mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

#### SEE INSIDE BOOKLET FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND COMPLETE DIRECTIONS FOR USE.

AD 100410

EPA Reg. No. 5905-196

EPA Est. No.: First letters of batch code indicate producing establishment: 5905-GA-1=CG, 5905-AR-1=WA, 5905-IA-1=DI, 5905-CA-1=KC

"RQ", Organophosphorus Pesticide, Liquid, Toxic (Malathion), UN3018

Manufactured For  
**HELENA CHEMICAL COMPANY**  
225 SCHILLING BOULEVARD, SUITE 300  
COLLIERVILLE, TENNESSEE 38017

### NET CONTENTS:

PEEL BACK BOOK HERE AND RESEAL AFTER OPENING

OPM #110221

# FYFANON®

## Malathion Insecticide

### ACTIVE INGREDIENT:

Malathion: O,O-dimethyl phosphorodithioate of diethyl mercaptosuccinate ..... 56.44%

**INERT INGREDIENTS** ..... 43.56%

**TOTAL** ..... 100.00%

(ONE GALLON OF FYFANON CONTAINS 5 POUNDS OF MALATHION)

### KEEP OUT OF REACH OF CHILDREN WARNING – AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you.)

**See Inside Booklet For Additional Precautions and Directions for Use.**

#### FIRST AID Organophosphate

**IF SWALLOWED:** Immediately call a poison control center or doctor. Do not induce vomiting unless told to by a poison control center or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. In case of emergency, call 1-800-424-9300.

**NOTE TO PHYSICIAN:** Contains an organophosphate that inhibits cholinesterase. Atropine is antidotal. 2PAM may be effective as an adjunct to atropine.

AD 100410

EPA Reg. No. 5905-196

EPA Est. No.: First letters of batch code indicate producing establishment:

5905-GA-1=CG, 5905-AR-1=WA, 5905-IA-1=DI, 5905-CA-1=KC

**“RQ”, Organophosphorus Pesticide, Liquid, Toxic (Malathion), UN3018**

Manufactured For

**HELENA CHEMICAL COMPANY**

225 SCHILLING BOULEVARD, SUITE 300 • COLLIERVILLE, TENNESSEE 38017 • PHONE: 901-761-0050

Job #58040

## PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING – AVISO

Causes substantial but temporary eye injury. Harmful if swallowed. Harmful if absorbed through skin. Harmful if inhaled. Do not get in eyes or on clothing. Avoid breathing vapor or spray mist.

### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are barrier laminate, butyl rubber, nitrile rubber, or viton. If you want more options, follow the instructions for category *F* on an EPA chemical-resistance category selection chart.

**For all formulations and all use patterns – mixers, loaders, applicators, flaggers and other handlers must wear:**

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves
- Chemical-resistant headgear (add for airblast application)
- Chemical-resistant apron (add for mixers and loaders)

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

### ENGINEERING CONTROLS

Pilots must use an enclosed cockpit in a manner that is consistent with the WPS for Agricultural Pesticides [40 CFR 170.240(d)(6)]. Pilots must wear the PPE required on this labeling for applicators.

### USER SAFETY RECOMMENDATIONS

#### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic organisms, including fish and invertebrates. This product may contaminate water through drift of spray in wind. This product has a high potential for runoff after application. Use care when applying in or to an area which is adjacent to any body of water and do not apply when weather conditions favor drift from target area. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product.

A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from rainfall runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

This pesticide is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI). The REI for each crop is listed in the directions for use associated with each crop.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Shoes plus socks, and
- Chemical-resistant gloves made of any waterproof material

## NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, or nurseries. Do not enter or allow others to enter until sprays have dried.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

**PESTICIDE STORAGE:** FYFANON® should be stored in the original unopened container in a secure dry place. Do not contaminate with other pesticides or fertilizers. The product should never be heated above 55°C (131°F), and should not be stored for long periods of time at a temperature in excess of 25°C (77°F).

**PESTICIDE DISPOSAL:** To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

**CONTAINER DISPOSAL: Nonrefillable container (equal to or less than 5 gallons):** Do not reuse or refill this container. Triple rinse (or equivalent). Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

**Nonrefillable container (greater than 5 gallons):** Do not reuse or refill this container. Triple rinse (or equivalent). Triple rinse as follows: Empty the remaining contents into application equipment or mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

## PRECAUTIONS AND RESTRICTIONS

### SPRAY DRIFT REQUIREMENTS

Observe the following requirements when spraying in the vicinity of aquatic areas such as, but not limited to lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish ponds.

### DROPLET SIZE

Use the largest droplet size consistent with acceptable efficacy. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible, and by avoiding excessive spray boom pressure.

For groundboom and aerial applications, use only medium or coarser spray nozzles according to ASAE (S572) definition for standard nozzles, or a volume mean diameter (VMD) of 300 microns or greater for spinning atomizer nozzles. In conditions of low humidity and high temperatures, applicators should use a coarser droplet size.

**WIND DIRECTION AND SPEED**

Make aerial or ground applications when the wind velocity favors on target product deposition (approximately 3 to 10 mph). Do not apply when wind velocity exceeds 15 mph. Avoid applications when wind gusts approach 15 mph. For all non-aerial applications, wind speed must be measured adjacent to the application site on the upwind side, immediately prior to application.

**TEMPERATURE INVERSION**

Do not make aerial or ground applications into areas of temperature inversions.

Inversions are characterized by stable air and increasing temperatures with increasing distance above the ground. Mist or fog may indicate the presence of an inversion in humid areas. Where permissible by local regulations, the applicator may detect the presence of an inversion by producing smoke and observing a smoke layer near the ground surface.

In conditions of low humidity and high temperatures, applicators should use a coarser droplet size.

**ADDITIONAL REQUIREMENTS FOR GROUND APPLICATIONS**

Spray should be released at the lowest height consistent with pest control and flight safety. Applications more than 10 feet above the crop canopy should be avoided. For groundboom applications, apply with nozzle height no more than 4 feet above the ground or crop canopy.

For airblast applications, turn off outward pointing nozzles at row ends and when spraying the outer two rows. To minimize spray loss over the top in orchard applications, spray must be directed into the canopy.

**BUFFER ZONES FOR AERIAL APPLICATION**

When making a Non-ULV application with aerial application equipment, a minimum buffer zone of 25 feet must be maintained along any water body.

**ADDITIONAL REQUIREMENTS FOR AERIAL APPLICATIONS**

For aerial applications, the spray boom should be mounted on the aircraft as to minimize drift caused by wingtip or rotor vortices. The minimum practical boom length should be used and must not exceed 75% of wing span or 90% rotor diameter.

Aerial applicators must consider flight speed and nozzle orientation in determining droplet size.

When applications are made with a cross-wind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

**AGRICULTURAL USES**

Use the higher rate when insect pressure is heavy.

Make application to agricultural sites using a minimum of 30 (ground) or 5 (aerial) gallons of water/A unless otherwise noted in the table directly below.

Crop	Pests Controlled	Rate/Acre	Max. Single App. Rate (lb ai/A)	Max. # of App. per year	Min. App. Interval (days)	Min. Pre-Harvest Interval (days)	Restricted Entry Interval (days)
Alfalfa	Alfalfa weevil larvae*; aphids; grasshoppers; lygus bugs; potato leaf hoppers; spider mites; spittlebugs; stink bugs; pea aphid	1.5-2 pints	1.25	2 per cutting	14	0	12 hrs
	Armyworms Clover leaf weevil Vetch bruchid	2 pints 1.5 pints 2 pints					
* Apply to blooming alfalfa only in early morning or evening when bees are not working field or not hanging on outside of hives. * Apply when day temperature is expected to be above 65°F and when 50-70% of leaves show damage.							

(continued)

Crop	Pests Controlled	Rate/Acre	Max. Single App. Rate (lb ai/A)	Max. # of App. per year	Min. App. Interval (days)	Min. Pre-Harvest Interval (days)	Restricted Entry Interval (days)
<b>Apricots</b>	Aphids; codling moth; European fruit lecanium; orange tortrix; soft brown scale; terrapin scale	2 pints	1.5	2	7	6	12 hrs
<b>Avocado</b>	Greenhouse thrips; latnia scale; omnivorous looper; orange tortrix; soft brown scale	7.5 pints	4.7	2	30	7	2 days
• Application rates are based on a standard dilution rate of 500 (ground) gallons of water/A.							
<b>Barley</b>	Cereal leaf beetle; English grain aphids; grasshoppers; greenbugs; winter grain mites	1-2 pints	1.25	2	7	7	12 hrs
<b>Beets, garden</b>	Aphids	1.5-2 pints	1.25	3	7	7	12 hrs
• Do not apply to Sugar Beets.							
<b>Blueberry (high bush and low bush)</b>	Blueberry maggots; cherry fruit worm; cranberry fruit worm; Japanese beetle	2 pints	1.25	3	5	1	12 hrs
• The rates for use on blueberries are based on a standard of 200 gallons per acre dilute spray.							
<b>Broccoli; Chinese Broccoli; Broccoli Raab</b>	Aphids; cabbage looper; imported cabbageworm; carrot weevil; flea beetle	1-2 pints	1.25	2	7	2	2 days

(continued)

Crop	Pests Controlled	Rate/Acre	Max. Single App. Rate (lb ai/A)	Max. # of App. per year	Min. App. Interval (days)	Min. Pre-Harvest Interval (days)	Restricted Entry Interval (days)
<b>Brussels sprouts</b>	Aphids; cabbage looper; imported cabbage worm; carrot weevil; flea beetle	1-2 pints	1.25	2	7	2	2 days
<b>Cabbage</b>	Aphids; cabbage looper; imported cabbageworm; diamondback moth; webworm; carrot weevil; flea beetle	1-2 pints	1.25	6	7	7	2 days
• For caterpillars on summer and fall plantings, begin when true leaves appear. On other plantings for listed insects, apply when first appear.							
<b>Chinese cabbage</b>	Aphids; cabbage looper; imported cabbageworm; diamondback moth; webworm; carrot weevil; flea beetle	1-2 pints	1.25	2	7	7	24 hours
• For caterpillars on summer and fall plantings, begin when true leaves appear. On other plantings for listed insects, apply when first appear.							
<b>Caneberries (blackberry; boysenberry; dewberry; gooseberry; loganberry; raspberry)</b>	Aphids; rose scale chafers; Japanese beetle; leafhoppers; mites; thrips	3 pints	2.0	3	7	1	12 hrs
• Application rates are based on a standard dilution rate of 200 (ground) gallons of water/A.							

(continued)

Crop	Pests Controlled	Rate/Acre	Max. Single App. Rate (lb ai/A)	Max. # of App. per year	Min. App. Interval (days)	Min. Pre-Harvest Interval (days)	Restricted Entry Interval (days)
Cucumber	Aphids; pickleworms; spider mites; cut worms; darkling ground beetle; leafhoppers; squash vine borer; thrips	1.5-2.8 pints	1.75	2	7	1	24 hrs
Celery	Aphids; spider mites	2.4 pints	1.5	2	7	7	24 hrs
Cherries (sweet and tart)	Black cherry aphid; fruit tree leafroller; Japanese beetle; cherry fruit fly; eyespotted bud moth	2.8 pints	1.75	4	3	3	12 hrs
• Application rates are based on a standard dilution rate of 400 (ground) gallons of water/A for mature trees. Injury may occur on sweet cherries, particularly in the Northwest.							
Citrus Fruits (grapefruit; lemon; lime; orange; tangerine; tangelo)	Thrips; California red scale; yellow scale; purple scale; black scale; soft scale; citricola scale	All states other than CA:					
		4.5 pints or 1.5 pints	4.5 or 1.5	1 or 3	N/A or 30	7	3 days or 12 hrs
		CA only:					
		7.5 pints or 1.5 pints	7.5 or 1.5	1 or 3	N/A or 30	7	3 days or 12 hours
• Use the higher rate for heavy infestation. • Apply with sufficient water to obtain full coverage of foliage. Apply in a minimum of 200 gallons of water per acre by ground, and up to 500 gallons of water per acre for mature trees. Use higher volumes of water as appropriate to ensure thorough coverage of foliage, depending on density and size of area to be treated.							

(continued)



Crop	Pests Controlled	Rate/Acre	Max. Single App. Rate (lb ai/A)	Max. # of App. per year	Min. App. Interval (days)	Min. Pre-Harvest Interval (days)	Restricted Entry Interval (days)
Clover	Alfalfa weevil larvae; aphids; grasshoppers; lygus bugs; potato leaf hoppers; spider mites; spittlebugs; stink bugs; pea aphid	1.5-2 pints	1.25	2 per cutting	14	0	12 hrs
	Armyworms Clover leaf weevil Vetch bruchid	2 pints 1.5 pints 2 pints					
• Do not apply to clover in bloom.							
Collards	Aphids; harlequin cabbage bug; smaller cabbage looper; leaf hoppers; leaf miners	1.5 pints	1.0	3	7	7	12 hrs
Corn (field)	Aphids; corn earworms; corn rootworm adults; young grasshoppers; sap beetle; thrips; smaller armyworms; leaf hopper	1.5 pints	1.0	2	7	7	3 days for detassling; 12 hrs for all other activities
Corn (sweet and pop)				2	5	5	3 days for detassling; 12 hrs for all other activities
• For corn earworm and sap beetles, treat when 10% of the ears show silk. • Injury may occur in the whorl silk stages.							

(continued)

Crop	Pests Controlled	Rate/Acre	Max. Single App. Rate (lb ai/A)	Max. # of App. per year	Min. App. Interval (days)	Min. Pre-Harvest Interval (days)	Restricted Entry Interval (days)
Cotton	Brown cotton leafworm; cotton aphid; cotton leaf-worm; cotton leaf perforator; desert spider mite; leafhoppers; lygus bugs; thrips; whiteflies; fall armyworms; garden webworms; grasshoppers	1.5-4 pints	2.5	3	7	7	2 days
	Boll weevil	2-4 pints					
	Cotton fleahoppers	1-1.5 pints					
	Lygus bugs; thrips	4 pints					
• Use higher rates for larger insects or heavy infestations.							
Dandelion	Aphids	1.5-2 pints	1.25	2	7	7	24 hrs
Eggplant	Aphids; spider mites	1.6-2.5 pints	1.56	4	5	3	12 hrs
Endive (escarole)	Aphids; spider mites	1.5-2 pints	1.25	2	7	7	24 hrs
Figs	Dried fruit beetle; vinegar flies	3 pints + 1-2 gal. sulfured molasses	2.0	2	5	5	24 hrs
Garlic	Aphids; thrips	1.5-2.5 pints	1.56	3	7	3	24 hrs

(continued)

Crop	Pests Controlled	Rate/Acre	Max. Single App. Rate (lb ai/A)	Max. # of App. per year	Min. App. Interval (days)	Min. Pre-Harvest Interval (days)	Restricted Entry Interval (days)
Grain Elevators & Silos (stored barley, corn, oats, rye, wheat)	Cereal leaf beetle; confused flour beetle; flat grain beetle; granary weevil; Indian meal moth; lesser grain borer; maize weevil; red flour beetle; rice weevil; rusty grain beetle; saw-toothed grain beetle	Mix 8 pints/25 gallons of water.	5 lb/25 gallons	1 per storage period	N/A	N/A	12 hrs
<ul style="list-style-type: none"><li>• Do not apply directly to grain.</li><li>• Before applying spray, clean thoroughly.</li><li>• Remove and burn all sweeping and debris.</li><li>• For a residual wall, floor and machinery spray in grain elevators and silos, before loading grain, make a thorough application.</li></ul>							
Grapes (raisin, table, wine)	Leafhoppers; spidermites; European fruit lecanium*; Drosophila; Japanese beetle; terrapin scale	3 pints	1.88	2	14	3	3 days for girdling and tying  24 hrs for all other activities
	Mealybugs	1.5 pints					
	Grape Vines (overwintering on root stock only)	Grape phylloxera					
<ul style="list-style-type: none"><li>• Application rates are based on a standard dilution rate of 200 (ground) gallons of water/A.</li><li>• Injury may occur on grapes of Almeria, Cardinal, Italia and Ribier varieties when sprays containing Fyfanon 57% EC are applied after clusters appear.</li><li>• Remove excess soil from roots and submerge entire root system in the Fyfanon solution for 5 minutes.</li><li>• Agitate solution at all times.</li></ul> <p>* Make full coverage applications when newly hatched nymphs are migrating over vines, usually shortly after bloom.</p>							

(continued)

Crop	Pests Controlled	Rate/Acre	Max. Single App. Rate (lb ai/A)	Max. # of App. per year	Min. App. Interval (days)	Min. Pre-Harvest Interval (days)	Restricted Entry Interval (days)
<b>Grass, forage, hay (Bermuda, barnyard grass, canary grass, fescue, orchardgrass, red top, timothy, yellow foxtail)</b>	Cereal leaf beetle; aphids; leafhoppers; grasshoppers	2 pints	1.25	1 per cutting	N/A	0	12 hrs
<b>Horseradish</b>	Aphids; diamondback moth; flea beetles; leafhoppers	1.5-2 pints	1.25	3	7	7	24 hrs
<b>Kale</b>	Aphids; cabbage looper; imported cabbageworm; webworm; diamondback moth	1.5 pints	1.0	3	5	7	12 hrs
<b>Kohlrabi</b>	Aphids; cabbage looper; imported cabbageworm; diamondback moth; webworm; carrot weevil; flea beetle	1-2 pints	1.25	2	7	7	24 hrs
<b>Kumquats</b>	Thrips; California red scale; yellow scale; purple scale; black scale; soft scale; citricola scale	4.5 pints	4.5	1	N/A	7	2 days

- Apply with sufficient water to obtain full coverage of foliage. Apply in a minimum of 200 gallons of water per acre by ground, and up to 500 gallons of water per acre for mature trees. Use higher volumes of water as appropriate to ensure thorough coverage of foliage, depending on density and size of area to be treated.

(continued)

Crop	Pests Controlled	Rate/Acre	Max. Single App. Rate (lb ai/A)	Max. # of App. per year	Min. App. Interval (days)	Min. Pre-Harvest Interval (days)	Restricted Entry Interval (days)
Leek	Aphids Onion maggot flies	1.5-2.5 pints	1.56	2	7	3	24 hrs
Lettuce (head)	Aphids; leaf-hoppers; spider mites; cabbage looper	3 pints	1.88	2	6	14	24 hrs
Lettuce (leaf)		2.5-3 pints	1.88	2	5	14	24 hrs
Mint	Aphids; flea beetles; leafhoppers; spider mites; caterpillars	1.5 pints	0.94	3	7	7	12 hrs
Mustards (mustard greens; mustard spinach; Chinese mustard mizuna)	Aphids; cabbage looper; imported cabbageworm; webworm; diamondback moth	1.5 pints	1.0	3	5	7	12 hrs
Nectarines	Spider mites; plum curculio	1-2 pints	3.0	3	7	7	24 hrs
	Aphids*; Japanese beetles*	4-4.8 pints					
• Application rates are based on a standard dilution rate of 300 (ground) gallons of water/A. • Fyfanon 57% EC may cause fruit spotting on nectarines. * May be mixed with spray oil for dormant and delayed dormant applications. Follow spray oil manufacturer's directions.							
Oats	Cereal leaf beetle; English grain aphids; young grasshoppers; greenbugs	1.5 pints	1.0	2	7	7	12 hrs
Okra	Aphids Japanese beetle	1.5 pints 1.9 pints	1.2	5	7	1	12 hrs
Onion (bulb and green)	Onion maggots	1.5-2.5 pints	1.56	2	7	3	12 hrs
	Onion thrips	1.5-2 pints					

(continued)

Crop	Pests Controlled	Rate/Acre	Max. Single App. Rate (lb ai/A)	Max. # of App. per year	Min. App. Interval (days)	Min. Pre-Harvest Interval (days)	Restricted Entry Interval (days)
Parsnip	Aphids; cabbage looper; imported cabbageworm; diamondback moth; webworm; carrot weevil; flea beetle	1.5-2 pints	1.25	3	7	7	24 hrs
Peaches	Black cherry aphid; black peach aphid; European red mite; green peach aphid; rusty plum aphid; Japanese beetle; spider mite	2.4 pints	3.0	3	11	7	24 hrs
	Oriental fruit moth; plum curculio; cottony peach scale; European fruit lecanium; ter-rapin scale	4.8 pints					
• Application rates are based on a standard dilution rate of 300 (ground) gallons of water/A. • Do not apply more than 9 pounds of actual Fyfanon 57% EC per acre per year to peach trees.							
Peas (dried, green)	Pea weevils; aphids	1.5 pints	1.0	2	7	3	12 hrs
Pecans	Pecan bud moth; aphids; pecan leaf case-bearer <sup>2</sup> ; spider mites; mites; Pecan phylloxera <sup>1</sup> ; Pecan nut casebearer <sup>2</sup> ; Walnut husk fly <sup>2</sup> ;	1.5-2 pints	2.5	2	7	7	24 hrs
• Application rates are based on a standard dilution rate of 500 (ground) gallons of water/A for mature trees 25-35 feet high. <sup>1</sup> Apply when buds begin to develop. <sup>2</sup> Apply when first generation eggs begin to hatch.							

(continued)

Crop	Pests Controlled	Rate/Acre	Max. Single App. Rate (lb ai/A)	Max. # of App. per year	Min. App. Interval (days)	Min. Pre-Harvest Interval (days)	Restricted Entry Interval (days)
<b>Peppers</b>	Aphids Pepper maggots	1-2.5 pints 2.5 pints	1.56	2	5	3	12 hrs
<b>Potatoes</b>	Aphids; grasshoppers; leafhoppers	2 pints	1.56	2	7	0	12 hrs
	False chinch bug	1.5 pints					
	Mealybugs	2-2.5 pints					
<b>Radish</b>	Aphids	1.5 pints	1.0	3	7	7	12 hrs
<b>Rutabagas</b>	Aphids	1.5 pints	1.0	3	7	7	12 hrs
<b>Rice Rice (wild)</b>	Rice stink bug; rice leaf miner	2 pints	1.25	2	7	7	12 hrs
<ul style="list-style-type: none"> <li>• Treat for leafminers shortly after first rice blades appear on surface of the water.</li> <li>• For leafminers, apply when the eggs and larvae are abundant on the seedling rice.</li> <li>• Apply during early milk and dough stage using a minimum of 2 (aerial) gallons of water/A.</li> <li>• Do not apply Propanil within 15 days of malathion treatment.</li> </ul> <p><b>NOTE FOR AQUATIC USES (rice):</b> Broadcast use only over intermittently flooded areas. Application may not be made around bodies of water where fish or shellfish are grown and/or harvested.</p>							
<b>Rye</b>	Cereal leaf beetle	1-1.5 pints	1.0	3	7	7	12 hrs
	English grain aphids; young grasshoppers; greenbugs	1.5 pints					
<b>Salsify</b>	Aphids; imported cabbage worm; cabbage looper; carrot weevil; flea beetles; leafhoppers; spider mites; thrips	1-2 pints	1.25	3	7	7	24 hrs
<b>Shallot</b>	Aphids; thrips	2.5 pints	1.56	2	7	3	24 hrs
<b>Sorghum</b>	Greenbugs	1.5 pints	1.0	2	7	7	12 hrs

(continued)

Crop	Pests Controlled	Rate/Acre	Max. Single App. Rate (lb ai/A)	Max. # of App. per year	Min. App. Interval (days)	Min. Pre-Harvest Interval (days)	Restricted Entry Interval (days)
<b>Spinach</b>	Aphids	1.5 pints	1.0	2	7	7	12 hrs
<b>Squash, summer</b>	Aphids; cucumber beetle; leaf miners; pickleworms; spider mites; cut worms; darkling ground beetle; leafhoppers; squash vine borer; thrips	2.8 pints	1.75	3	7	1	24 hrs
<b>Squash, winter</b>	Aphids	1.5 pints	1.0	3	7	1	12 hrs
<b>Strawberry</b>	Aphids; spider mites; Field crickets; lygus bugs; spittle bugs; thrips; potato leafhopper; strawberry leafroller; strawberry root weevil; white flies; thrips	1.5-3.2 pints	2.0	4	7	3	12 hrs
<b>Sweet Potatoes</b>	Leafhoppers; Morningglory leaf miner	1.5-2.5 pints	1.56	2	7	0	12 hrs
<b>Swiss chard</b>	Aphids	1.5 pints	1.0	2	7	14	12 hrs
<b>Tomatoes, Tomatillos</b>	Aphids; spider mites Drosophila	1.5 pints 2.5 pints	1.56	4	5	1	12 hrs
• Apply a full coverage application to fruit and foliage.							

(continued)



Crop	Pests Controlled	Rate/Acre	Max. Single App. Rate (lb ai/A)	Max. # of App. per year	Min. App. Interval (days)	Min. Pre-Harvest Interval (days)	Restricted Entry Interval (days)
<b>Vetch</b>	Alfalfa weevil larvae; aphids; armyworms; clover leaf weevil; grasshoppers; lygus bugs; pea aphid; potato leafhoppers; spider mites; spittlebugs; vetch bruchid; omnivorous leaf tier; pea aphid; vetch bruchid	1-2 pints	1.25	2 per cutting	14	0	12 hrs
<b>Watercress</b>	Aphids	2 pints or 1.6 pints	1.25 or 1.0	5	3	3	24 hrs or 12 hrs
<b>Wheat (Spring and Winter)</b>	Cereal leaf beetle	1-1.5 pints	1.0	2	7	7	12 hrs
	English grain aphids; young grasshoppers; greenbugs	1.5 pints					

#### NON-AGRICULTURAL OUTDOOR USE SITES

Site	FL. OZ./Acre	Max. Single App. Rate	Use Pattern Limitations	Restricted Entry Interval (days)
<b>Christmas tree plantations</b>	82	3.2 lb ai/A	Maximum of 2 applications per year	12 hrs
<b>Fence rows/hedge rows</b>	6	0.2439 lb ai/1,000 ft <sup>2</sup>		
<b>Ornamental and/or shade trees</b>	64	2.5 lbs ai/100 gal	Maximum of 2 applications per year; 10-day minimum re-treatment interval	12 hrs
<b>Ornamental herbaceous plants</b>	64	2.5 lbs ai/100 gal		12 hrs
<b>Ornamental non-flowering plants</b>	64	2.5 lbs ai/100 gal		

(continued)

<b>Ornamental woody shrubs and vines</b>	64	2.5 lbs ai/100 gal	Maximum of 2 applications per year/growing cycle; 10-day minimum re-treatment interval	12 hrs
<b>Pine seed orchards</b>	82	3.2 lbs ai/A	Maximum of 2 applications per year/growing season; 7-day minimum re-treatment interval	12 hrs

#### OUTDOOR FLY CONTROL

**Fly control:** For use on the lower outside foundation of the home and fence/hedge rows.

PEST CONTROLLED	RATE	DIRECTIONS FOR USE
<b>Adult flies</b>	Straight sprays: 5 tablespoons + 1 gallon water or 1 cup + 2-1/2 gallons water or 6 ounces + 12 gallons water	Apply as a spray at the rate of 1 gallon per 1,000 sq. ft. on painted surfaces and 2 gallons per 1,000 sq. ft. on unpainted surfaces where flies alight or congregate
<b>Adult flies Fly maggots</b>	Bait sprays: 5 tablespoons + 7 tablespoons sugar or molasses (unsulfurized) or corn syrup + 2-1/2 gallons water or 1 cup + 1 cup sugar or molasses (unsulfurized) or corn syrup + 2-1/2 gallons water or 1 quart + 2-1/2 lbs. sugar or 1 quart molasses (unsulfurized) or 1 quart corn syrup + 12 gallons water	Apply as a bait spray. Do not apply to freshly whitewashed surfaces. Wait 14 days after whitewashing before applying.
Repeat applications as necessary. Avoid applying oil-based formulations to valuable ornamental plants as injury may occur.		

#### MOSQUITO CONTROL (Localized Applications only – not for wide areas of applications)

**OUTDOOR RESIDENTIAL ADULT MOSQUITO CONTROL:** For control of mosquitoes outdoors, use a 2% to 5% Malathion area or fog spray. For a 2% spray, dilute 1 part of **FYFANON®** with 28 parts of water; for 5%, dilute 1 part of **FYFANON®** with 11 parts of water. Repeat as necessary. Treat along lower outside foundation of home, and spot treat shrubbery and vegetation where mosquitoes may rest.

**MOSQUITO LARVAE CONTROL:** For control of mosquito larvae in standing water, intermittently flooded areas, stagnant water, and temporary rain pools, apply 13 ounces of **FYFANON®** per acre. Mix in sufficient water or oil to obtain even coverage when applied by air or ground equipment. NOTE: BROADCAST USE ONLY OVER INTERMITTENTLY FLOODED AREAS. APPLICATION MAY NOT BE MADE AROUND BODIES OF WATER WHERE FISH OR SHELLFISH ARE GROWN AND/OR HARVESTED COMMERCIALY. CONTAMINATION OF SHALLOW, FISH-BEARING WATERS MAY KILL FISH.

## **CONDITIONS OF SALE – LIMITED WARRANTY AND LIMITATIONS OF LIABILITY AND REMEDIES**

**Read the Conditions of Sale – Warranty and Limitations of Liability and Remedies before using this product. If the terms are not acceptable, return the product, unopened, and the full purchase price will be refunded.**

The directions on this label are believed to be reliable and must be followed carefully. Insufficient control of pests and/or injury to the crop to which the product is applied may result from the occurrence of extraordinary or unusual weather conditions or the failure to follow the label directions or good application practices, all of which are beyond the control of Helena Chemical Company (the "Company") or seller. In addition, failure to follow label directions may cause injury to crops, animals, man or the environment. The Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the directions for use subject to the factors noted above which are beyond the control of the Company. To the extent consistent with applicable law, the Company makes no other warranties or representations of any kind, express or implied, concerning the product, including no implied warranty of merchantability or fitness for any particular purpose, and no such warranty shall be implied by law.

The exclusive remedy against the Company for any cause of action relating to the handling or use of this product shall be limited to, at Helena Chemical Company's election, one of the following:

1. Refund of the purchase price paid by buyer or user for product bought, or
2. Replacement of the product used

To the extent consistent with applicable law, the Company shall not be liable and any and all claims against the Company are waived for special, indirect, incidental, or consequential damages or expense of any nature, including, but not limited to, loss of profits or income. The Company and the seller offer this product and the buyer and user accept it, subject to the foregoing conditions of sale and limitation of warranty, liability and remedies.

© Copyright Helena Holding Company, 2010.

**FYFANON®** is a registered trademark of Cheminova.



