

Pro

LEBANON

Scape®

HOMOGENEOUS
FERTILIZER
CONFRONT® + TEAM®
ANNUAL GRASS AND BROADLEAF WEED CONTROL

19-0-7

0 8 8 6 8 5 5 4 3 5 0 0

F699

21-54350

GUARANTEED ANALYSIS

Total Nitrogen (N) 19%
8.0% Ammoniacal Nitrogen
5.8% Water Insoluble Nitrogen*
2.6% Urea Nitrogen
2.6% Other Water Soluble Nitrogen*
Soluble Potash (K₂O) 7%
Iron (Fe) 0.5%
0.5% Water Soluble Iron (Fe)
Derived from: Ammonium Phosphate, Ammonium Sulfate, Methylene Ureas, Urea, Muriate of Potash,
Ferric Oxide, Ferrous Sulfate.
Chlorine (Cl) not more than 9.0%
*8.4% Slowly Available Nitrogen from Methylene Ureas.
NOTICE: This product contains the secondary nutrient iron. Iron may stain concrete surfaces and should not be applied on dry
or water dampened concrete and should be removed from these areas promptly after application by sweeping or blowing. Do
not wash off with water.

SUGGESTED SPREADER SETTINGS

Spreader	Settings 3.6#/1000 sq. ft	Spreaders	Settings 3.6#/1000 sq. ft
LebanonTurf	4½	PrizeLAWN BF & CBR III	K
PennMulch HVO	K	PrizeLAWN CBR II	4
ProScape SS	K	PrizeLAWN LF-1	3½
Cyclone	4½	Republic Rotary	4½
Earthway Rotary	15	Scotts AccuPro	K
Gandy	26	Scotts (Rotary) R7X	H
Lely (HR, 1250,1500,2010)	5II	Scotts (Rotary) R8A	L
Lesco (letter dial/numeric) I/17	I/17	Spyker	4½
Ortho Rotary	4½	Vicon (all models)	24

The above spreader settings are approximate. These settings were calibrated and field tested.
However, age and condition of spreader, speed of operation, and evenness of terrain may require
slightly different settings for desired coverage.

ACTIVE INGREDIENTS:

Triethylamine salt of triclopyr 0.50%
Triethylamine salt of clopyralid. 0.18%
Benefin 0.82%
Trifluralin. 0.43%
OTHER INGREDIENTS: 98.07%
TOTAL 100.00%

Acid Equivalent: triclopyr - 0.36%
clopyralid - 0.12%

Triclopyr CAS No. 57213-69-1
Clopyralid CAS No. 1702-17-6

Benefin CAS No. 1861-40-1
Trifluralin CAS No. 1582-09-8
(L) (D)
EPA Reg. No. 961-391
EPA Est. No. 961-PA-01:961-IL-01

CONFRONT and TEAM are registered trademarks of Dow AgroSciences LLC. Superscript used is first letter of lot number.
ProScape is a registered trademark of Lebanon Seaboard Corporation.

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID

IF ON SKIN
OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

IF SWALLOWED:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

IF INHALED:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
- Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For
Emergency Medical Information call (888) 208-1368.

See back panel for additional Precautionary Statements

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT.
USE ONLY ACCORDING TO LABEL INSTRUCTIONS.

Before buying this product, read the Warranty Section elsewhere on this label.
If terms are not acceptable, return the unopened container to the place of purchase at once.

For technical assistance or more information about our products visit www.LebanonTurf.com
Manufactured by:
Lebanon Seaboard Corporation
1600 E. Cumberland St. • Lebanon, PA 17042
Made in the U.S.A.
www.LebanonTurf.com

NET WEIGHT

50

LBS. (22.68 KG)

HOMOGENEOUS FERTILIZER CONFRONT® + TEAM® ANNUAL GRASS AND BROADLEAF WEED CONTROL

19-0-7



F699

21-54350

GUARANTEED ANALYSIS

Total Nitrogen (N)	19%
8.0% Ammoniacal Nitrogen	
5.8% Water Insoluble Nitrogen*	
2.6% Urea Nitrogen	
2.6% Other Water Soluble Nitrogen*	
Soluble Potash (K₂O)	7%
Iron (Fe)	0.5%
0.5% Water Soluble Iron (Fe)	

Derived from: Ammonium Phosphate, Ammonium Sulfate, Methylene Ureas, Urea, Muriate of Potash, Ferric Oxide, Ferrous Sulfate.

Chlorine (Cl) not more than 9.0%

*8.4% Slowly Available Nitrogen from Methylene Ureas.

NOTICE: This product contains the secondary nutrient iron. Iron may stain concrete surfaces and should not be applied on dry or water dampened concrete and should be removed from these areas promptly after application by sweeping or blowing. Do not wash off with water.

SUGGESTED SPREADER SETTINGS

Settings		Settings	
Spreader	3.6#/1000 sq. ft	Spreaders	3.6#/1000 sq. ft
LebanonTurf	4½	PrizeLAWN BF & CBR III	K
PennMulch HVO	K	PrizeLAWN CBR II	4
ProScape SS	K	PrizeLAWN LF-1	3½
Cyclone	4½	Republic Rotary	4½
Earthway Rotary	15	Scotts AccuPro	K
Gandy	26	Scotts (Rotary) R7X	H
Lely (HR, 1250,1500, 2010)	5II	Scotts (Rotary) R8A	L
Lesco (letter dial/numeric) I/17		Spyker	4½
Ortho Rotary	4½	Vicon (all models)	24

The above spreader settings are approximate. These settings were calibrated and field tested. However, age and condition of spreader, speed of operation, and evenness of terrain may require slightly different settings for desired coverage.

Triethylamine salt of triclopyr	0.50%
Triethylamine salt of clopyralid	0.18%
Benefin	0.82%
Trifluralin	0.43%
OTHER INGREDIENTS:	<u>98.07%</u>
TOTAL	100.00%

ProScape is a registered trademark of Lebanon Seaboard Corporation.

FIRST AID

OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

IF SWALLOWED:

- Call a poison control center or doctor for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

IF INHALED:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
- Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For Emergency Medical Information call (888) 208-1368.

See back panel for additional Precautionary Statements

READ THE ENTIRE LABEL BEFORE USING THIS PRODUCT.

USE ONLY ACCORDING TO LABEL INSTRUCTIONS.

Before buying this product, read the Warranty Section elsewhere on this label.

If terms are not acceptable, return the unopened container to the place of purchase at once.



For technical assistance or more information about our products visit www.LebanonTurf.com

Manufactured by:

Lebanon Seaboard Corporation

1600 E. Cumberland St. • Lebanon, PA 17042

Made in the U.S.A.

www.LebanonTurf.com



FERTILIZER WITH CONFRONT® + TEAM® ANNUAL GRASS AND BROADLEAF WEED CONTROL

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Personal Protective Equipment (PPE):

All loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of any waterproof material

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to freshwater, marine, and estuarine fish and aquatic invertebrates including shrimp and oyster. Do not apply in a manner that will directly expose canals, lakes, streams, ponds, marshes or estuaries to aerial drift. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organism in water adjacent to treated areas.

Clopyralid is a chemical that which can travel (seep or leach) through soil and under certain conditions contaminate ground water which may be used for irrigation or drinking purposes. Users are advised not to apply clopyralid where soils have a rapid to very rapid permeability throughout the profile (such as loamy sand to sand) and the water table of an over limestone bedrock, severely fractures surfaces, and substrates which could allow direct introduction into an aquifer. Your local agencies can provide further information on the type of soil in your area and the location of ground water. Triclopyr has properties and characteristics associated with chemicals detected in groundwater. The use of triclopyr in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all directions carefully before applying this product.

Not for use on plants grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

Application Restrictions: Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during applications. Maximum of two applications per year.

Entry Restrictions: This product must be watered in as soon as possible after application. Do not enter or allow people (or pets) to enter the treated area until dusts have settled. Except for those people involved in the watering-in - do not enter or allow other people (or pets) to enter the treated areas until the watering-in is complete and the surface is dry.

GENERAL INFORMATION

This product provides broad-spectrum control of broadleaf weeds and annual grasses in established cool season or warm season turfgrasses. Confront herbicide is particularly effective against established broadleaf weeds such as dandelions and Team herbicide prevents the establishment of annual grass weeds, including crabgrass. This product is not effective against established grass weeds.

This product may be used on the following turf species:

Established Cool Season Turfgrasses - Kentucky bluegrass, chewings fescue, creeping red fescue, sheeps fescues, tall fescue, perennial ryegrass.

Established Warm Season Turfgrasses⁺ - Bahiagrass (Argentine or Pensacola), Common Bermudagrass, Buffalograss, Centipedegrass, Zoysiagrass (*Zoysia japonica*, *Zoysia tenuifolia*), Tall Fescue (growing in warm season area).

⁺Do not treat warm season turfgrasses with this product when mowing height is less than 1/2 inch. Do not use this product on golf course putting greens and tees.

PRECAUTIONS AND RESTRICTIONS

Do not treat warm season turfgrasses with this product when mowing height is less than 1/2 inch. Do not use this product on golf course putting greens and tees.

PRECAUTIONS AND RESTRICTIONS

FOR USE ONLY ON NONRESIDENTIAL TURF SUCH AS ATHLETIC AND RECREATIONAL SPORTS FIELDS, CEMETERIES, GOLF COURSES, INDUSTRIAL SITES, NON-CROPLAND, PARKS, RIGHTS -OF-WAY AND ROADSIDES.

MAXIMUM OF TWO APPLICATIONS PER YEAR.

DO NOT USE ON RESIDENTIAL TURF.

Do not apply this product to bentgrass; hybrid bermudagrass varieties such as Tifway, Tifdwarf, and Tifgreen; St. Augustinegrass; and ground covers such as lippia.

This product may discolor and/or stunt turf that is not well established or is stressed or weakened due to unfavorable climatic conditions, temperature extremes, drought, nematodes, or other factors which damage or weaken turf. Apply only to healthy, well-established turf that has a well-anchored root system.

Common bermudagrass, centipedegrass, zoysiagrass, and bahiagrass may be temporarily discolored by this product.

On bahiagrass and centipedegrass, do not apply before mid-March or until after the second mowing of the year. In the fall, do not treat centipedegrass after the first of October or bahiagrass after the first of November.

Do not apply this product to newly seeded turf until turfgrass is fully established. Do not seed, sod, or sprig after application. Do not reseed until 12 weeks after application. Delay sodding or sprigging until 4 weeks after application.

Do not collect grass clippings for mulch or compost. Do not send clippings to a compost facility. Applicator must give notice to landowners/property managers to not use grass clippings for composting. Do not graze treated areas or feed clippings to livestock.

Do not apply this product directly to, or otherwise permit it to come into direct contact with vegetables, flowers, fruit or ornamental trees, cotton, grapes, tobacco, or other nontarget broadleaf plants. Do not apply when weather conditions favor drift from target areas. Drift or runoff may adversely affect nontarget plants.

Do not apply this product to exposed roots of shallow rooted trees and shrubs or allow contact with exposed suckers or injury may occur. Do not apply at greater than specified rates, especially under the branch spread (dripline) of trees.

WEEDS CONTROLLED

When applied as directed, this product will control the following weeds:

Annual Grass Weeds: Barnyardgrass, Crabgrass (smooth and hairy), Goosegrass, Foxtail, Annual Bluegrass (*Poa annua*).

Established Broadleaf Weeds: Black Medic, Bristly Hawksbeard, Broadleaf Plantain, Burdock, Canada Thistle, Catsear (Coast Dandelion), Coffee Weed, Common Chickweed, Common Cocklebur, Common Dandelion, Common Ragweed, Common Vetch, Common Yellow Woodsorrel, Creeping Beggarweed, Creeping Woodsorrel, Curly Dock, Dollarweed, Dwarf Beggarweed, English Lawn Daisy, False Dandelion, Goldenrod, Ground Ivy, Hawkweed, Henbit, Hop Clover, Lambsquarters, Lespedeza, Matchweed, Mouseear Chickweed, Musk Thistle, Narrow Leaf Plantain (Buckhorn), Old World Diamond Flower, Poison Ivy, Prostrate Spurge, Red Clover, Roundleaf Mallow, Sheep Sorrel, Shepherdspurse, Slender Speedwell, Smartweed, Spurweed, Virginia Buttonweed, Virginia Pepperweed, White Clover, Wild Buckwheat, Wild Violet.

Note: Moist foliage application will provide quicker results on broadleaf weeds. For best herbicide effect when applying to dry foliage, apply when good soil moisture conditions are present. A repeat application may be needed for harder to control weeds in 4 to 6 weeks.

APPLICATION DIRECTIONS HOW TO APPLY THIS PRODUCT

Apply this product at a rate of 3.6 lbs. per 1,000 sq. ft. (157 lbs./acre). This application rate is equivalent to 0.75 lb. ai/acre of Confront and 2 lbs. ai/acre of Team. Make application using equipment that will ensure uniform coverage. For postemergence control of broadleaves, application should be made when weeds are actively growing.

Good soil moisture conditions should be present before application to achieve optimal results.

Application under drought conditions may provide less than desirable results.

Broadleaf weed species germinate at different times. Emerged broadleaf weeds present at time of application are controlled by Confront, unemerged grassy weeds present at time of application are controlled by Team.

Note: If a second application is made, do not apply less than 4 weeks apart to minimize the potential for grass injury. Newly seeded turf should be mowed 4 times before being treated.

Note: Supplemental applications of lawn fertilizer are needed to achieve a minimum 1 pound nitrogen per 1,000 sq. ft.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store product in its original container in a cool, dry, locked place out of the reach of children and animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Completely empty container into application equipment.

Then dispose of empty container in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

WARRANTY

The Manufacturer warrants that this product conforms to its chemical description and is reasonably fit for the purpose on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to Manufacturer and to the extent consistent with applicable law, Buyer assumes the risks of any such use.

MANUFACTURED BY:
LEBANON SEABOARD CORPORATION

1600 EAST CUMBERLAND STREET

LEBANON, PA 17042

For technical assistance or more information about our products visit www.LebanonTurf.com

Information regarding the contents and levels of metals in this product is available
on the Internet at <http://www.regulatory-info-lebsea.com>